

Symmetrical Peripheral Gangrene: An Unusual Occurrence in a 1-Month Old: A Case Report

Authors: I. K. Ndu¹; O. C. Nduagubam¹; C. C. Iheji^{1*}; I. N. Asinobi¹; U. Ekwochi¹

Affiliations: ¹Department of Pediatrics, Enugu State University Teaching Hospital, Parklane

ABSTRACT

INTRODUCTION: Symmetrical peripheral gangrene (SPG) is a disfiguring vascular disorder described as symmetrical distal ischemic damage leading to gangrene of two or more sites in the absence of large vessel obstruction or vasculitis. Although it tends to occur in all age groups, the youngest recorded age to our knowledge is 3.5 months. The etiologies are many and have been grouped into infectious and non-infectious causes. Bacteria and viruses such as *Staphylococcus aureus* and HIV and low flow states, as seen in hypovolemic shock for example, have been implicated.

CASE PRESENTATION: The case we describe is a 4-week-old, HIV-exposed, small for age (SGA) preterm female who presented with passage of mucoid stool, non-projectile vomiting and continuous fever. At presentation, she was in respiratory distress and was febrile, and appeared pale with features of moderate dehydration. She had hepatomegaly and hypoactive bowel sounds. Blood culture yielded moderate growth of *Staphylococcus aureus*. Appropriate antibiotics were commenced. Within 20 hours of admission, she developed dry gangrene which started as bluish discoloration of the right hand progressing to the lower third of the forearm, and within a few hours her left hand and subsequently both feet were involved. Her clinical condition further deteriorated at which point she was noticed to have progressive hypoxia and a small volume pulse despite all interventions. She died before planned amputation.

DISCUSSION: This case report underscores the possibility of SPG in neonates despite its rarity and the need for clinicians to have high index of suspicion.

Keywords: Symmetrical Peripheral Gangrene, Neonate, Immunosuppression, *Staphylococcus Aureus*, Case Report

INTRODUCTION

Symmetrical peripheral gangrene (SPG) is a well-documented, disfiguring vascular disorder complicating diverse medical conditions. It has been described as symmetrical distal ischemic damage leading to gangrene of two or more sites in the absence of large vessel obstruction or vasculitis [1-3]. The exact incidence is not known and patients of any age can be affected [1]. However, the disease remains even more rare in children and the youngest age recorded to our knowledge is 3.5 months [5,6].

The etiologies of SPG are many and could be grouped into infectious and non-infectious causes [1]. The infectious causes include bacteria (*Streptococcus pneumoniae*, *Staphylococcus aureus*, *Neisseria meningitidis*, *Streptococcus pyogenes*, *Enterococcus faecalis*), protozoans (*Plasmodium falciparum*), and viruses (HIV, Rubeola virus, and Varicella zoster virus). Non-infectious causes include cardiovascular pathologies (hypertension, heart failure, myocardial infarction, hypovolemic shock, supraventricular tachycardia, pulmonary embolism), drugs (dopamine, adrenaline, noradrenaline), malignancy (acute lymphoblastic leukemia, Hodgkin's lymphoma), connective tissue disorders (systemic lupus erythematosus, antiphospholipid antibodies, polymyalgia rheumatic, progressive systemic sclerosis, Henoch-Schonlein purpura, Takayasu arteritis), miscellaneous (dog bites, appendicitis, extra corporeal shock wave lithotripsy, suprapubic prostatectomy, cholecystectomy, sickle cell disease, hyperosmolar coma, hypernatremic dehydration), protein C or S or antithrombin III deficiency [1,7-17].

Though many pathophysiological agents have been implicated,

Though many pathophysiological agents have been implicated,

***Corresponding author:** Iheji Chukwunonso Chigozie, Department of Paediatrics, Enugu State University Teaching Hospital, Parklane; E-mail: gozieprimenet@yahoo.com; Tel: +2347033806550; **Potential Conflicts of Interest (CoI):** All authors: no potential conflicts of interest disclosed; **Funding:** All authors: no funding was disclosed; **Academic Integrity:** All authors confirm that they have made substantial academic contributions to this manuscript as defined by the ICMJE; **Ethics of human subject participation:** The study was approved by the local Institutional Review Board. Informed consent was sought and gained where applicable; **Originality:** All authors: this manuscript is original has not been published elsewhere; **Type-editor:** Ahmed (USA) **Review:** This manuscript was peer-reviewed by three reviewers in a double-blind review process; **Received:** 11th February 2019; **Initial decision given:** 3rd July 2019; **Revised manuscript received:** 15th September 2019; **Accepted:** 10th October 2019 **Copyright:** © The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND) (click here) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Biomedical Centre (RBC)/Rwanda Health Communication Center, P.O.Box 4586, Kigali. **ISSN:** 2079-097X (print); 2410-8626 (online) **Citation for this article:** I. K. Ndu; O.C. Nduagubam; C.C. Iheji. et al. Peripheral Gangrene: An Unusual Occurrence in a 1-Month Old – A Case Report. Rwanda Medical Journal, Vol. 77, no. 1, pp. 35-38, 2020.

the exact mechanism by which they cause vascular occlusion is largely unknown [1]. Low flow state or septicemia are almost invariably present. Fever followed by marked coldness, pallor, cyanosis, pain, restricted mobility of extremity should always raise suspicion of SPG [18]. The investigations to determine the etiology may not be able to identify the exact cause, but the clinical presentation and the suspected etiologies should guide the choice of investigations. SPG is associated with high rates of morbidity and mortality. Amputation rates reported range from 30 to 50% and mortality has been shown to be as high as 40% [13,19,20].

We present a case that is particularly notable for the very young age of the patient and the rarity of the condition. This patient's course highlights the need for clinicians to have a high index of suspicion because early intervention with antibiotics, intravenous fluids, judicious use of inotropes, anticoagulants, and reduction or removal of aggravating factors is essential line of treatment [1].

OUR CASE

A four-week-old female presented to the Children's Emergency Room (CHER) of the Enugu State University Teaching Hospital, Parklane, Enugu, Enugu State, Nigeria, with poor weight gain which started from birth, passage of mucoid stool that began one week prior to presentation, non-projectile vomiting and continuous fever that commenced a day prior to presentation. She was delivered at home at 36-weeks gestational age by a single mother whose antenatal visits were not kept and she had previously been unaware of her HIV-positive status. The pregnancy however was said to be devoid of any adverse event until at 36-weeks gestation when the mother spontaneously ruptured membrane and was delivered of a live female neonate who cried at birth with a birth weight of 2.2kg (9th centile). The child had inadequate nutritional intake from birth as she was not exclusively breastfed and was introduced to pap from birth.

At presentation, she was in respiratory distress and febrile (38.6°C). She was moderately pale with features of moderate dehydration. Her pulse was 142bpm, regular, and full volume. Her respiratory rate was 52 breaths per minute. Her weight was 2.4kg (0.4th centile). She was noted to have hepatomegaly of 4cm and hypoactive bowel sounds.

The admitting diagnosis was sepsis with moderate dehydration on a background of malnutrition. The HIV screening was reactive/positive. A complete blood count showed marked leukocytosis of $23.04 \times 10^9/L$ of which 76% were neutrophils with many band forms; and blood culture demonstrated moderate growth of *Staphylococcus aureus* sensitive to ceftriaxone, levofloxacin, cefixime and vancomycin. She was commenced on intranasal oxygen, and intravenous ceftriaxone and vancomycin.

About 20-hours into admission, she was noticed to have bluish discoloration of the right hand and her pulse volume was also noted to be reduced. Dopamine was commenced at $5\mu g/kg/minute$ and the intravenous fluid volume was readjusted.

A few hours later, the bluish discoloration of the right hand progressed to the lower third of the forearm. At this point, dopamine was discontinued, intravenous metronidazole and vancomycin were added and the child was being worked up for surgery. Within 48-hours both feet up to lower third of the legs and left hand up to the lower third of the forearm were noticed to have been involved [Figure 1]. However, she died before the planned surgical intervention the following day.



Figure 1: Gangrene involving all four limbs

DISCUSSION

This case describes a neonate with symmetrical peripheral gangrene involving all limbs and her case was striking for the very young age at which it occurred, compared to the youngest case previously recorded in the literature at an age of 3.5 months [5,6].

This patient did have several risk factors associated with SPG. First, an immature immune system is well recognized to be present in neonates, especially preterm infants as in this case. Decreased function of neutrophils and other cells involved in the response to infection has been demonstrated in both term and preterm infants. Preterm infants may also have low concentrations of immunoglobulins. Both preterm and term infants have quantitative and qualitative defects of the complement system [21]. In addition, the underweight and malnourished status of our patient may have also contributed to the immunosuppression.

The relationship between malnutrition and increased susceptibility to and mortality from infectious diseases is well documented in children. Decreased gastric acid secretion promotes bacterial overgrowth and this, coupled with mucosal atrophy and the consequent hypolactasia, cause diarrhea and malabsorption [21,22]. Moreover, there is extensive atrophy of tissues including the thymus, spleen, and other lymphoid tissues, resulting in reduction of circulating lymphocytes. Both humoral and cell-mediated immunity, including lymphokine production, are compromised in malnutrition. While phagocytic function may be normal in such children,

monocyte chemotaxis as well as intracellular killing is depressed. Levels of most components of the complement system are also reduced. Our patient had diarrhea which may be related to the malnutrition, though mucosal atrophy was not demonstrated histologically [22]. HIV infection is also well known for its immunosuppressive effect. HIV has been noted to evade the immune system by causing depletion of certain subset of killer T-Lymphocytes, which further become dysfunctional. Also early in the disease, patients lose HIV-specific CD4+ T-cell responses including the secretion of interferons that normally slow the replication of viruses. PCR was not performed to confirm the diagnosis of HIV in the patient because of her death prior to completion of investigations and treatments.

Our patient's blood culture yielded moderate growth of *S. aureus*. This is consistent with the several case reports of SPG [29, 6-8,18].

Third, use of dopamine has also been implicated in SPG [10,11]. This may not have been the cause of the SPG in the index case as the gangrene had already started in the first affected limb before it

REFERENCES

- SK Ghosh , D Bandyopadhyay , "Symmetrical peripheral gangrene," Indian J Dermatol Venereol Leprol [Internet]. vol. 77, no. 2, pp. 244 – 8, 2011. Accessed on: Nov. 6, 2018 [Online]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/21393969>
- AH Kalajian , KB Turpen , KO Donovan , JC Malone , JP Calen , "Phenylephrine-Induced Microvascular Occlusion Syndrome in a Patient With a Heterozygous Factor V Leiden Mutation," Arch Dermatol [Internet]. vol. 143, no. 10, pp. 1314 – 7, Oct.1, 2007. Accessed on: Nov. 6, 2018 [Online]. Available:<http://www.ncbi.nlm.nih.gov/pubmed/17938347>
- P Chittichai , N Chierakul, MT Davis , "Peripheral Gangrene in Nonfatal Pediatric Cerebral Malaria," Southeast Asian J Trop Med Public Health. vol. 22, no.2, June 1991.
- F Ejagwulu, A Ibrahim, "Symmetrical peripheral gangrene and extremity amputations," Ann Niger Med [Internet]. vol. 8, no. 2, pp. 98, 2014 [cited 2018 Nov 7];8(2):98. Available: <http://www.anmjournals.com/text.asp?2014/8/2/98/153363>
- JC Urimubabo , F Byiringiro, R Ssebuufu, V Fawcett , "Case Report Symmetrical Peripheral gangrene in two children at the University Teaching Hospital of Butare (CHUB): a report of two cases". RMJ . vol. 72, no. 3, pp. 25 – 7, Sept., 2015.
- S Aytac , G Kanra , G Secmeer , "Outcome in Children With Purpura Fulminans : Report on 16 Patients," American Journal of Hematology. vol. 80, 2005.
- MDP Davis , KM Dy , S Nelson , "Presentation and outcome of purpura fulminans associated with peripheral gangrene in 12 patients at Mayo Clinic," J Am Acad Dermatol [Internet]. vol. 57, no. 6, pp. 944 – 56, Dec. 1, 2007. Accessed on: Nov. 6, 2018. [Online]. Available: <http://linkinghub.elsevier.com/retrieve/pii/S0190962207011954>
- S Ghosh , D Bandyopadhyay, A Ghosh , "Symmetrical peripheral gangrene: a prospective study of 14 consecutive cases in a tertiary-care hospital in eastern India," J Eur Acad Dermatology Venereol [Internet]. vol. 24, no. 2, pp. 214 – 8, Feb. 2010. Accessed on: Nov. 7 2018. [Online]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/19522719>
- S Tripathy, B Rath, "Symmetric peripheral gangrene: Catch it early!" J Emerg Trauma Shock [Internet]. vol. 3, no. 2, pp. 189 – 90, Apr., 2010. Accessed on: Dec. 15, 2018 [Online]. Available:<http://www.ncbi.nlm.nih.gov/pubmed/20606800>
- RC McGouran , GA Emmerson, "Symmetrical peripheral gangrene," Br Heart J [Internet]. vol. 39, no. 5, pp. 569 – 72, May 1, 1977. Accessed on: Nov. 6, 2018 [Online]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/861101>
- R Kashyap, RK Behl, S Mahajan, P Jaret, RK Patial, SS Kaushal, "Symmetrical peripheral gangrene due to viral gastroenteritis," J Assoc Physicians India [Internet]. vol. 52, pp. 500 – 1, Jun. 2004. Accessed on: Nov. 7, 2018 [Online]. Available:<http://www.ncbi.nlm.nih.gov/pubmed/15645964>
- J Pedraz, Y Delgado-Jiménez, A González-De Arriba, L Ríos-Buceta, J Fernández-Herrera, A García-Diez, "Peripheral symmetrical gangrene," Clin Exp Dermatol [Internet]. vol. 34, no. 4, pp. 552 – 4, Jun. 2009. Accessed on: Dec. 15, 2018 [Online]. Available:<http://www.ncbi.nlm.nih.gov/pubmed/19522990>
- MA Molos , JC Hall, "Symmetrical peripheral gangrene and disseminated intravascular coagulation," Arch Dermatol [Internet]. vol. 121, no. 8, pp. 1057 – 61, Aug. 1985. Accessed on: Nov. 6, 2018 [Online]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/3896158>
- JH Yoo , JK Min , SS Kwon , CH Jeong , WS Shin , "Symmetrical peripheral gangrene complicating Klebsiella pneumoniae sepsis associated with antiphospholipid antibodies," Ann Rheum Dis [Internet]. vol. 63, no. 4, pp. 459 – 60. Apr. 2004. Accessed on: Nov. 6, 2018 [Online]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/15020345>
- SJ Mellor, NJ Cheshire, "Symmetrical peripheral gangrene after extracorporeal shockwave lithotripsy" BJU Int [Internet]. vol. 84, no. 1, pp. 167 – 8. Jul. 1999. Accessed on: Nov. 8, 2018 [Online]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/10444142>

was commenced and it was also quickly discontinued. However, dopamine may have contributed to the rapid progression of the clinical course.

Finally, other possible triggers such as protein C, protein S or antithrombin III deficiency were not assayed, though clinically the baby did not demonstrate any bedside features of bleeding disorder.

CONCLUSION

We recommend that physicians attending to infants should have a high index of suspicion of the early manifestation of SPG once cyanosis of an extremity is noticed. The role of early surgical intervention could play an important role in preventing the mortality associated with SPG.

Consent for publication

Written consent for the report was obtained from the mother.

16. A Crisci, G Giannarini, C Selli, "Symmetrical Peripheral Gangrene after Suprapubic Prostatectomy," *Scand J Urol Nephrol* [Internet]. vol. 36, no. 6, pp. 478 – 80. Jan. 9, 2002. Accessed on: Nov. 7, 2018 [Online] Available: <http://www.ncbi.nlm.nih.gov/pubmed/12623515>
17. SK Ghosh, B Majumder, S Ghosh, S Chatterjee, M Agarwal, "Symmetrical peripheral gangrene complicating ventricular pseudoaneurysm: a report of an unusual case and a brief review of the literature," *An Bras Dermatol* [Internet] vol. 91, no. 5 Supl 12016;91(5 Supl 1):S169-71.
18. S Tiwary, R Shankar, R Khanna, A Khanna, "Symmetrical Peripheral Gangrene," *The Internet Journal of Surgery*. vol. 7, no 2, Apr. 2005.
19. EA Kirsch, RP Barton, L Kitchen, BP Giroir, "Pathophysiology, treatment and outcome of meningococemia: a review and recent experience," *Pediatr Infect Dis J* [Internet]. vol. 15, no. 11, pp. 967 – 78, Nov, 1996. Accessed on: Nov. 9, 2018 [Online]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/8933544>
20. MC Genoff, MM Hoffer, B Achauer, P Formosa, "Extremity amputations in meningococemia-induced purpura fulminans," *Plast Reconstr Surg* [Internet]. vol. 89, no. 5, pp. 878 – 81, May 1992. Accessed on: Nov. 9, 2018 [Online]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/1561259>
21. BJ Stoll, "Infections of the Neonatal Infant" in *Nelson's Textbook of Pediatrics 19th ed.*, R.M. Kliegman, B.F. Stanton, J.W. St. Geme III, N.F. Schor, and R.E. Behrman, Eds. Philadelphia: Elsevier Saunders, 2011, ch. 103, pp. 629 – 48.
22. TO Ulas and J. Ebenebe, "Nutritional Disorders in Childhood" in *Paediatrics and Child Health in a Tropical Region 2nd ed.*, J.C. Azubuike and K.E.O. Nkanginieme, Eds. Owerri, Nigeria: African Education Services, 2007, ch. 33, pp. 250 – 267.
23. A U Eneh, "Human Immune Deficiency Virus (HIV) Infection" in *Paediatrics and Child Health in a Tropical Region 2nd ed.*, J.C. Azubuike and K.E.O. Nkanginieme, Eds. Owerri, Nigeria: African Education Services, 2007, ch. 73, pp. 643 – 654.
24. NA Okoko, KO Owuor, JL Kulzer, GO Owino, IA Ogolla, RW Wandera, et al, "Factors associated with mother to child transmission of HIV despite overall low transmission rates in HIV-exposed infants in rural Kenya," *Int J STD AIDS* [Internet]. vol. 28, no. 12, pp. 1215 – 23, Oct. 9, 2017. Accessed on: Jan. 9, 2019 [Online]. Available: <http://journals.sagepub.com/doi/10.1177/0956462417693735>
25. S Ngwende, NT Gombe, S Midzi, M Tshimanga, G Shambira, A Chadambuka, "Factors associated with HIV infection among children born to mothers on the prevention of mother to child transmission programme at Chitungwiza Hospital, Zimbabwe, 2008," *BMC Public Health* [Internet]. vol. 13, pp. 1181, Dec. 14, 2013. Accessed on: [Jan. 5, 2019]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/24330311>
26. HIV & AIDS Information :: How likely is mother-to-child transmission? - What factors increase the chance of transmission from mother to baby? [Internet]. [cited 2019 Jan 3]. Available from: <http://www.aidsmap.com/What-factors-increase-the-chance-of-transmission-from-mother-to-baby/page/1321400/>
27. GS Birkhead, WP Pulver, BL Warren, S Hackel, D Rodríguez, L Smith, "Acquiring human immunodeficiency virus during pregnancy and mother-to-child transmission in New York: 2002-2006," *Obstet Gynecol* [Internet]. vol. 115, no. 6, pp. 1247 – 55, Jun. 2010. Accessed on: Jan. 3, 2019. [Online]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/20502297>
28. SH Landesman, LA Kalish, DN Burns, H Minkoff, HE Fox, C Zorrilla, et al, "Obstetrical Factors and the Transmission of Human Immunodeficiency Virus Type 1 from Mother to Child," *N Engl J Med* [Internet]. vol. 334, no. 25, pp. 1617 – 23, Jun. 20, 1996. Accessed on: Dec. 16, 2018 [Online]. Available: <http://www.nejm.org/doi/abs/10.1056/NEJM199606203342501>
29. SP Rai, LS Neog, D Bhattacharyya, M Ganguli, "Symmetrical Peripheral Gangrene complicating Staphylococcal Toxic Shock Syndrome". *MJAFI* 2008; 64 : 181-182